1.4 Product Information

1.4.1 Prescribing Information

Summary of Product Characteristics

1. NAME OF THE MEDICINAL PRODUCT

MONAST 10 (Montelukast Sodium Tablets 10 mg)

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Beige, rounded square shaped film coated tablets debossed with “I” on one side and ‘114’ on the other side

3. PHARMACEUTICAL FORM

Film-coated tablet

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Montelukast 10 mg film-coated tablets is indicated in the treatment of asthma as add-on therapy in those patients with mild to moderate persistent asthma who are inadequately controlled on inhaled corticosteroids and in whom “as-needed” short acting beta-agonists provide inadequate clinical control of asthma. In those asthmatic patients in whom Montelukast 10 mg film-coated tablets is indicated in asthma, Montelukast 10 mg film-coated tablets can also provide symptomatic relief of seasonal allergic rhinitis.

Montelukast 10 mg film-coated tablets in the prophylaxis of asthma in which the predominant component is exercise-induced bronchoconstriction.

4.2 Posology and method of administration

Posology:

The recommended dose for adults and adolescents 15 years of age and older with asthma, or with asthma and concomitant seasonal allergic rhinitis is one 10 mg tablet daily to be taken in the evening.

General recommendations:

The therapeutic effect of Montelukast 10 mg film-coated tablets on parameters of asthma control occurs within one day. Montelukast 10 mg film-coated tablets may be taken with or without food. Patients should be advised to continue taking Montelukast 10 mg film-coated tablets even if their asthma is under control, as well as during periods of worsening asthma. Montelukast 10 mg film-coated tablets should not be used concomitantly with other products containing the same active ingredient, montelukast.

No dosage adjustment is necessary for the elderly, or for patients with renal insufficiency, or mild to moderate hepatic impairment. There are no data on patients with severe hepatic impairment. The dosage is the same for both male and female patients.

Therapy with Montelukast 10 mg film-coated tablets in relation to other treatments for asthma

Montelukast 10 mg film-coated tablets can be added to a patient’s existing treatment regimen.

Inhaled corticosteroids:

Treatments with Montelukast 10 mg film-coated tablets can be used as add-on therapy in patients when inhaled corticosteroids plus "as needed" short acting β-agonists provide inadequate clinical control. Montelukast 10 mg film-coated tablets should not be abruptly substituted for inhaled corticosteroids

Paediatric population

Do not give Montelukast 10 mg film-coated tablets to children less than 15 years of age. The safety and efficacy of Montelukast 10 mg film-coated tablets in children less than 15 years has not been established.

Other available strengths/pharmaceutical forms:

5 mg chewable tablets are available for paediatric patients 6 to 14 years of age.

4 mg chewable tablets are available for paediatric patients 2 to 5 years of age.

4 mg granules are available for paediatric patients 6 months to 5 years of age.

Method of administration:

Oral use.

4.3 Contraindications

Hypersensitivity to the active substance(s) or to any of the excipients listed in section 6.1.

4.4 Special warnings and precautions for use

Patients should be advised never to use oral montelukast to treat acute asthma attacks and to keep their usual appropriate rescue medication for this purpose readily available. If an acute attack occurs, a short-acting inhaled beta-agonist should be used. Patients should seek their doctor's advice as soon as possible if they need more inhalations of short-acting beta-agonists than usual.

Montelukast should not be substituted abruptly for inhaled or oral corticosteroids.

There are no data demonstrating that oral corticosteroids can be reduced when montelukast is given concomitantly.

In rare cases, patients on therapy with anti-asthma agents including montelukast may present with systemic eosinophilia, sometimes presenting with clinical features of vasculitis consistent with Churg-Strauss syndrome, a condition which is often treated with systemic corticosteroid therapy. These cases have been sometimes associated with the reduction or

withdrawal of oral corticosteroid therapy. Although a causal relationship with leukotriene receptor antagonism has not been established, physicians should be alert to eosinophilia, vasculitis rash, worsening pulmonary symptoms, cardiac complications, and/or neuropathy presenting in their patients. Patients who develop these symptoms should be reassessed and their treatment regimens evaluated.

Treatment with montelukast does not alter the need for patients with aspirin-sensitive asthma to avoid taking aspirin and other non-steroidal anti-inflammatory drugs.

Patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicine.

Neuropsychiatric events have been reported in adults, adolescents, and children taking Montelukast 10 mg film-coated tablets . Patients and physicians should be alert for neuropsychiatric events. Patients and/or caregivers should be instructed to notify their physician if these changes occur. Prescribers should carefully evaluate the risks and benefits of continuing treatment with Montelukast 10 mg film-coated tablets if such events occur.

4.5 Interaction with other medicinal products and other forms of interaction

Montelukast may be administered with other therapies routinely used in the prophylaxis and chronic treatment of asthma. In drug-interactions studies, the recommended clinical dose of montelukast did not have clinically important effects on the pharmacokinetics of the following medicinal products: theophylline, prednisone, prednisolone, oral contraceptives (ethinyl estradiol/ norethindrone 35/1), terfenadine, digoxin and warfarin.

The area under the plasma concentration curve (AUC) for montelukast was decreased approximately 40% in subjects with co-administration of Phenobarbital. Since montelukast is metabolized by CYP 3A4, 2C8, and 2C9, caution should be exercised, particularly in children, when montelukast is co-administered with inducers of CYP 3A4, 2C8, and 2C9, such as phenytoin, Phenobarbital and rifampicin.

In vitro studies have shown that montelukast is a potent inhibitor of CYP 2C8. However, data from a clinical drug-drug interaction study involving montelukast and rosiglitazone (a probe

substrate representative of medicinal products primarily metabolized by CYP 2C8) demonstrated that montelukast does not inhibit CYP 2C8 in vivo. Therefore, montelukast is not anticipated to markedly alter the metabolism of medicinal products metabolized by this enzyme (e.g., paclitaxel, rosiglitazone, and repaglinide.)

In vitro studies have shown that montelukast is a substrate of CYP 2C8, and to a less significant extent, of 2C9, and 3A4. In a clinical drug-drug interaction study involving montelukast and gemfibrozil (an inhibitor of both CYP 2C8 and 2C9) gemfibrozil increased the systemic exposure of montelukast by 4.4-fold. No routine dosage adjustment of montelukast is required upon co-administration with gemfibrozil or other potent inhibitors of CYP 2C8, but the physician should be aware of the potential for an increase in adverse reactions.

Based on in vitro data, clinically important drug interactions with less potent inhibitors of CYP 2C8 (e.g., trimethoprim) are not anticipated. Co-administration of montelukast with itraconazole, a strong inhibitor of CYP 3A4, resulted in no significant increase in the systemic exposure of montelukast.

4.6 Fertility, pregnancy and lactation

Use during pregnancy

Animal studies do not indicate harmful effects with respect to effects on pregnancy or embryonal/foetal development.

Available data from published prospective and retrospective cohort studies with montelukast use in pregnant women evaluating major birth defects have not established a drug-associated risk. Available studies have methodologic limitations, including small sample size, in some cases retrospective data collection, and inconsistent comparator groups.

Montelukast 10 mg film-coated tablets may be used during pregnancy only if it is considered to be clearly essential.

Use during breastfeeding

Studies in rats have shown that montelukast is excreted in milk. It is unknown whether montelukast/metabolites are excreted in human milk.

Montelukast 10 mg film-coated tablets may be used in breast-feeding only if it is considered to be clearly essential

4.7 Effects on ability to drive and use machines

Montelukast has no or negligible influence on the ability to drive and use machines. However, individuals have reported drowsiness or dizziness.

4.8 Undesirable effects

The frequency using the following convention: Common (≥1/100 to <1/10); Uncommon (≥1/1,000 to <1/100); Uncommon (≥1/1,000 to <1/100); Rare (≥1/10,000 to <1/1,000); Very rare (<1/10,000); not known (cannot be estimated from the available data).

Montelukast has been evaluated in clinical studies as follows:

- 10 mg film-coated tablets in approximately 4000 adult and adolescent_asthmatic patients 15 years of age and older.
- 10 mg film-coated tablets in approximately 400 adult and adolescent asthmatic patients with seasonal allergic rhinitis 15 years of age and older.
- 5 mg chewable tablets in approximately 1750 paediatric asthmatic patients 6 to 14 years of age.

The following drug-related adverse reactions in clinical studies were reported commonly (1/100 to <1/10) in asthmatic patients treated with montelukast and at a greater incidence than in patients treated with placebo:

Body system Class	**Adult Patients** **15 years and older** **(two 12-week studies; n=795)**	**Paediatric and adolescents Patients** **6 to 14 years old** **(one 8-week study; n=201)**

		(two 56-week studies; n=615)
Nervous system disorders	headache	headache
Gastrointestinal disorders	abdominal pain	

With prolonged treatment in clinical trials with a limited number of patients for up to 2 years for adults, and up to 12 months for paediatric and adolescents patients 6 to 14 years of age, the safety profile did not change.

Tabulated list of Adverse Reactions

Adverse reactions reported in post-marketing use are listed, by System Organ Class and specific Adverse Reactions, in the table below. Frequency Categories were estimated based on relevant clinical trials.

system Organ Class	**Adverse Experience Term**	**Frequency Category***
Infections and infestations	upper respiratory infection†	Very Common
Blood and lymphatic system disorders	increased bleeding tendency	Rare
Immune system disorder	hypersensitivity reactions including anaphylaxis	Uncommon
	hepatic eosinophilic infiltration	Very Rare
Psychiatric disorders	dream abnormalities including nightmares, insomnia, somnambulism, irritability, anxiety, restlessness, agitation including aggressive behaviour or hostility, depression, psychomotor hyperactivity (including irritability, restlessness, tremor§)	Uncommon

	disturbance in attention, memory impairment	Rare
	hallucinations, disorientation, suicidal thinking and behaviour (suicidality)	Very Rare
Nervous system disorder	dizziness, drowsiness paraesthesia/hypoesthesia, seizure	Uncommon
Cardiac disorders	palpitations	Rare
Respiratory, thoracic and mediastinal disorders	epistaxis	Uncommon
	Churg-Strauss Syndrome (CSS) (see section 4.4), pulmonary eosinophilia	Very Rare
Gastrointestinal disorders	diarrhoea‡, nausea‡, vomiting‡	Common
	dry mouth, dyspepsia	Uncommon
Hepatobiliary disorders	elevated levels of serum transaminases (ALT, AST)	Common
	hepatitis (including cholestatic, hepatocellular, and mixed-pattern liver injury).	Very Rare
Skin and subcutaneous tissue disorders	rash‡	Common
	bruising, urticaria, pruritus	Uncommon
	angiooedema	Rare
	erythema nodosum, erythema multiforme	Very Rare
Musculoskeletal, connective tissue and bone disorders	arthralgia, myalgia including muscle cramps	Uncommon
General disorders and	pyrexia‡	Common

administration site conditions	asthenia/fatigue, malaise, oedema,	Uncommon

*Frequency Category: Defined for each Adverse Experience Term by the incidence reported in the clinical trials data base: Very Common (≥1/10), Common (≥1/100 to </10), Uncommon (≥1/1000 to </100), Rare (≥1/10,000 to </1000), Very Rare (</10,000).

†This adverse experience, reported as Very Common in the patients who received montelukast, was also reported as Very Common in the patients who received placebo in clinical trials.

‡This adverse experience, reported as Common in the patients who received montelukast, was also reported as Common in the patients who received placebo in clinical trials.

§ Frequency Category: Rare

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via HPRA Pharmacovigilance. Website: www.hpra.ie. By reporting side effects you can help provide more information on the safety of this medicine.

4.9 Overdose

In chronic asthma studies, montelukast has been administered at doses up to 200 mg/day to patients for 22 weeks and in short term studies, up to 900 mg/day to patients for approximately one week without clinically important adverse experiences.

There have been reports of acute overdose in post-marketing experience and clinical studies with montelukast. These include reports in adults and children with a dose as high as 1,000 mg (approximately 61 mg/kg in a 42 month old child). The clinical and laboratory findings

observed were consistent with the safety profile in adults and paediatric patients. There were no adverse experiences in the majority of overdose reports.

Symptoms of overdose

There were no adverse experiences in the majority of overdose reports. The most frequently occurring adverse experiences were consistent with the safety profile of montelukast and included abdominal pain, somnolence, thirst, headache, vomiting and psychomotor hyperactivity.

Management of overdose

No specific information is available on the treatment of overdose with montelukast. It is not known whether montelukast is dialysable by peritoneal- or haemo-dialysis.

5. Pharmacological properties

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Other systemic drugs for obstructive airway diseases, Leukotriene receptor antagonist

ATC code: R03D C03

Mechanism of action

The cysteinyl leukotrienes (LTC_4, LTD_4, and LTE4) are potent inflammatory eicosanoids released from various cells including mast cells and eosinophils. These important pro-asthmatic mediators bind to cysteinyl leukotriene (CysLT) receptors. The CysLT type-1 ($CysLT_1$) receptor is found in the human airway (including airway smooth muscle cells and airway macrophages) and on other pro-inflammatory cells (including eosinophils and certain myeloid stem cells). CysLTs have been correlated with the pathophysiology of asthma and allergic rhinitis. In asthma, leukotriene-mediated effects include bronchoconstriction, mucous secretion, vascular permeability, and eosinophil recruitment. In allergic rhinitis, CysLTs are released from the nasal mucosa after allergen exposure during both early- and late-phase

reactions and are associated with symptoms of allergic rhinitis. Intranasal challenge with CysLTs has been shown to increase nasal airway resistance and symptoms of nasal obstruction.

Pharmacodynamic effects

Montelukast is an orally active compound which binds with high affinity and selectivity to the $CysLT_1$ receptor. In clinical studies, montelukast inhibits bronchoconstriction due to inhaled LTD_4 at doses as low as 5 mg. Bronchodilation was observed within 2 hours of oral administration. The bronchodilation effect caused by a beta agonist was additive to that caused by montelukast. Treatment with montelukast inhibited both early- and late phase bronchoconstriction due to antigen challenge. Montelukast, compared with placebo, decreased peripheral blood eosinophils in adult and paediatric patients. In a separate study, treatment with montelukast significantly decreased eosinophils in the airways (as measured in sputum) and in peripheral blood while improving clinical asthma control.

In studies in adults, montelukast, 10 mg once daily, compared with placebo, demonstrated significant improvements in morning FEV_1 (10.4% vs 2.7% change from baseline), AM peak expiratory flow rate (PEFR) (24.5 L/min vs 3.3 L/min change from baseline), and significant decrease in total beta-agonist use (-26.1% vs -4.6% change from baseline). Improvement in patient-reported daytime and nighttime asthma symptoms scores was significantly better than placebo.

Clinical efficacy and safety

In studies in adults, montelukast, 10 mg once daily, compared with placebo, demonstrated significant improvements in morning FEV1 (10.4% vs 2.7% change from baseline), AM peak expiratory flow rate (PEFR) (24.5 L/min vs 3.3 L/min change from baseline), and significant decrease in total β-agonist use (-26.1% vs -4.6% change from baseline). Improvement in patient-reported daytime and nighttime asthma symptoms scores was significantly better than placebo.

Studies in adults demonstrated the ability of montelukast to add to the clinical effect of inhaled corticosteroid (% change from baseline for inhaled beclometasone plus montelukast vs beclometasone, respectively for FEV_1: 5.43% vs 1.04%; beta-agonist use: -8.70% vs 2.64%). Compared with inhaled beclometasone (200 µg twice daily with a spacer device), montelukast demonstrated a more rapid initial response, although over the 12-week study, beclometasone provided a greater average treatment effect (% change from baseline for montelukast vs beclometasone, respectively for FEV_1: 7.49% vs 13.3%; beta agonist use: -28.28% vs -43.89%). However, compared with beclometasone, a high percentage of patients treated with montelukast achieved similar clinical responses (e.g., 50% of patients treated with beclometasone achieved an improvement in FEV_1 of approximately 11% or more over baseline while approximately 42% of patients treated with montelukast achieved the same response).

A clinical study was conducted to evaluate montelukast for the symptomatic treatment of seasonal allergic rhinitis in adult asthmatic patients 15 years of age and older with concomitant seasonal allergic rhinitis. In this study, montelukast 10 mg tablets administered once daily demonstrated a statistically significant improvement in the Daily Rhinitis Symptoms score, compared with placebo. The Daily Rhinitis Symptoms score is the average of the Daytime Nasal Symptoms score (mean of nasal congestion, rhinorrhea, sneezing, nasal itching) and the Nighttime Symptoms score (mean of nasal congestion upon awakening, difficulty going to sleep, and nighttime awakenings scores). Global evaluations of allergic rhinitis by patients and physicians were significantly improved, compared with placebo. The evaluation of asthma efficacy was not a primary objective in this study.

In an 8-week study in paediatric patients 6 to 14 years of age, montelukast 5 mg once daily, compared with placebo, significantly improved respiratory function (FEV_1 8.71% vs 4.16% change from baseline; AM PEFR 27.9 L/min vs 17.8 L/min change from baseline) and decreased "as-needed" beta-agonist use (-11.7% vs +8.2% change from baseline).

Significant reduction of exercise-induced broncho constriction (EIB) was demonstrated in a 12-week study in adults (maximal fall in FEV_1 22.33% for montelukast vs 32.40% for placebo; time to recovery to within 5% of baseline FEV_1 44.22 min vs 60.64 min). This

effect was consistent throughout the 12-week study period. Reduction in EIB was also demonstrated in a short term study in paediatric patients (maximal fall in FEV_1 18.27% vs 26.11%; time to recovery to within 5% of baseline FEV_1 17.76 min vs 27.98 min). The effect in both studies was demonstrated at the end of the once-daily dosing interval.

In aspirin-sensitive asthmatic patients receiving concomitant inhaled and/or oral corticosteroids, treatment with montelukast, compared with placebo, resulted in significant improvement in asthma control (FEV_1 8.55% vs -1.74% change from baseline and decrease in total beta-agonist use -27.78% vs 2.09% change from baseline).

5.2 Pharmacokinetic properties

Absorption

Montelukast is rapidly absorbed following oral administration. For the 10 mg film-coated tablet, the mean peak plasma concentration (Cmax) is achieved 3 hours (Tmax) after administration in adults in the fasted state. The mean oral bioavailability is 64%. The oral bioavailability and Cmax are not influenced by a standard meal. Safety and efficacy were demonstrated in clinical trials where the 10 mg film-coated tablet was administered without regard to the timing of food ingestion.

For the 5 mg chewable tablet, the Cmax is achieved in 2 hours after administration in adults in the fasted state. The mean oral bioavailability is 73% and is decreased to 63% by a standard meal.

Distribution

Montelukast is more than 99% bound to plasma proteins. The steady-state volume of distribution of montelukast averages 8-11 liters. Studies in rats with radiolabelled montelukast indicate minimal distribution across the blood-brain barrier. In addition, concentrations of radiolabelled material at 24 hours post-dose were minimal in all other tissues.

Biotransformation

Montelukast is extensively metabolised. In studies with therapeutic doses, plasma concentrations of metabolites of montelukast are undetectable at steady state in adults and children.

Cytochrome P450 2C8 is the major enzyme in the metabolism of montelukast. Additionally CYP 3A4 and 2C9 may have a minor contribution, although itraconazole, an inhibitor of CYP 3A4, was shown not to change pharmacokinetic variables of montelukast in healthy subjects that received 10 mg montelukast daily. Based on *in vitro* results in human liver microsomes, therapeutic plasma concentrations of montelukast do not inhibit cytochromes P450 3A4, 2C9, 1A2, 2A6, 2C19, or 2D6. The contribution of metabolites to the therapeutic effect of montelukast is minimal.

Elimination

The plasma clearance of montelukast averages 45 ml/min in healthy adults. Following an oral dose of radio labelled montelukast, 86% of the radioactivity was recovered in 5-day faecal collections and <0.2% was recovered in urine. Coupled with estimates of montelukast oral bioavailability, this indicates that montelukast and its metabolites are excreted almost exclusively via the bile.

Characteristics in patients

No dosage adjustment is necessary for the elderly or mild to moderate hepatic insufficiency. Studies in patients with renal impairment have not been undertaken. Because montelukast and its metabolites are eliminated by the biliary route, no dose adjustment is anticipated to be necessary in patients with renal impairment. There are no data on the pharmacokinetics of montelukast in patients with severe hepatic insufficiency (Child-Pugh score>9).

With high doses of montelukast (20- and 60-fold the recommended adult dose), decrease in plasma theophylline concentration was observed. This effect was not seen at the recommended dose of 10 mg once daily.

5.3 Preclinical safety data

In animal toxicity studies, minor serum biochemical alterations in ALT, glucose, phosphorus and triglycerides were observed which were transient in nature. The signs of toxicity in animals were increased excretion of saliva, gastro-intestinal symptoms, loose stools and ion imbalance. These occurred at dosages which provided>17-fold the systemic exposure seen at the clinical dosage. In monkeys, the adverse effects appeared at doses from 150 mg/kg/day (>232-fold the systemic exposure seen at the clinical dose). In animal studies, montelukast did not affect fertility or reproductive performance at systemic exposure exceeding the clinical systemic exposure by greater than 24-fold. A slight decrease in pup body weight was noted in the female fertility study in rats at 200 mg/kg/day (>69fold the clinical systemic exposure). In studies in rabbits, a higher incidence of incomplete ossification, compared with concurrent control animals, was seen at systemic exposure>24-fold the clinical systemic exposure seen at the clinical dose. No abnormalities were seen in rats. Montelukast has been shown to cross the placental barrier and is excreted in breast milk of animals.

No deaths occurred following a single oral administration of montelukast sodium at doses up to 5000 mg/kg in mice and rats (15,000 mg/m2 and 30,000 mg/m2 in mice and rats, respectively), the maximum dose tested. This dose is equivalent to 25,000 times the recommended daily adult human dose (based on an adult patient weight of 50 kg).

Montelukast was determined not to be phototoxic in mice for UVA, UVB or visible light spectra at doses up to 500 mg/kg/day (approximately>200-fold based on systemic exposure).

Montelukast was neither mutagenic in in vitro and in vivo tests nor tumorigenic in rodent species.

6. Pharmaceutical particulars

6.1 List of excipients

Lactose monohydrate Ph.Eur, Mannitol Ph.Eur, Croscarmellose sodium Ph.Eur, Hydroxypropyl cellulose Ph.Eur, Cellulose Microcrystallline Ph.Eur, magnesium stearate Ph.Eur, purified water.

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

24 months

6.4 Special precautions for storage

Store below 30˚C, protect from moisture.

6.5 Nature and contents of container

10's Blister (ALU-ALU)

6.6 Special precautions for disposal and other handling

Keep out of reach of children.

7. MARKETING AUTHORISATION HOLDER

HETERO LABS LIMITED (Unit-l),

Village: Kalyanpur, Chakkan Road,

Tehsil : Baddi, Distt.: Solan,

Himachal Pradesh -173 205, INDIA

Name : M/s Hetero Labs Limited, Unit V

Address : TSIIC Formulation SEZ, S.No 439, 440,441 & 458,

Polepally Village, Jadcherla

(Mandal), Mahaboob Nagar (District)-509301,

Telangana State, India.

7.2 Name and Address of Principal

Name : Hetero Labs Limited

Business Address : 7-2-A2, Hetero Corporate,
Industrial Estates, Sanath Nagar,
Hyderabad-500 018
Telangana.

Country	: INDIA
Telephone	: 0091-040-23704923/24

8. REGISTRATION NUMBER

Not applicable

9. DATE OF PUBLICATION OF THIS PACKAGE INSERT

Not applicable